

RESEARCH LETTER

10.1002/2016GL071772

Special Section:

The Arctic: An AGU Joint Special Collection

Key Points:

- Top-down and bottom-up estimates of the Arctic CH₄ natural sources do not match; we suggest that double-counting of emissions in bottom-up accounting is a major cause of this discrepancy
- Double-counting can occur due to overlap of wetland and lake or fluvial areas in models or inventories using low-resolution maps
- Double-counting can occur due to overlap of the δ¹³C-CH₄ property between many Arctic CH₄ sources

Supporting Information:

- Supporting Information S1

Correspondence to:

B. F. Thornton,
brett.thornton@geo.su.se

Citation:

Thornton, B. F., M. Wik, and P. M. Crill (2016), Double-counting challenges the accuracy of high-latitude methane inventories, *Geophys. Res. Lett.*, *43*, 12,569–12,577, doi:10.1002/2016GL071772.

Received 29 OCT 2016

Accepted 29 NOV 2016

Accepted article online 4 DEC 2016

Published online 29 DEC 2016

©2016. The Authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Double-counting challenges the accuracy of high-latitude methane inventories

Brett F. Thornton¹ , Martin Wik¹, and Patrick M. Crill¹ ¹Department of Geological Sciences and Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

Abstract Quantification of the present and future contribution to atmospheric methane (CH₄) from lakes, wetlands, fluvial systems, and, potentially, coastal waters remains an important unfinished task for balancing the global CH₄ budget. Discriminating between these sources is crucial, especially across climate-sensitive Arctic and subarctic landscapes and waters. Yet basic underlying uncertainties remain, in such areas as total wetland area and definitions of wetlands, which can lead to conflation of wetlands and small ponds in regional studies. We discuss how in situ sampling choices, remote sensing limitations, and isotopic signature overlaps can lead to unintentional double-counting of CH₄ emissions and propose that this double-counting can explain a pan-Arctic bottom-up estimate from published sources, 59.7 Tg yr⁻¹ (range 36.9–89.4 Tg yr⁻¹) greatly exceeding the most recent top-down inverse modeled estimate of the pan-Arctic CH₄ budget (23 ± 5 Tg yr⁻¹).

1. Introduction

At first glance, balancing the CH₄ budget should be simple. Decades long records of atmospheric CH₄ exist, and with a lifetime of less than 10 years in the atmosphere [Prather *et al.*, 2012], we should be able to construct a box model where known CH₄ sources minus known CH₄ sinks equals the current atmospheric burden. Though the atmospheric burden is well known, detailed accounting of both sources and sinks remains a tremendous challenge [Kirschke *et al.*, 2013], somewhat due to potentially large CH₄ sources newly noted in the past 10 years. Many of these potential new sources are regional, and many lie in the Arctic where warming temperatures may be more favorable for production and release of CH₄ from long-stored permafrost carbon (C), potentially contributing to a permafrost C warming feedback [Schuur *et al.*, 2015; Vonk *et al.*, 2013]. It remains a goal to reconcile the top-down Arctic CH₄ budget (e.g., calculating backward from the amount of CH₄ observed in the atmosphere to sources), with bottom-up budgets (e.g., summing the CH₄ sources and sinks to determine the atmospheric burden). We provide here an updated, but rough, bottom-up inventory, based on published estimates of various categories of natural Arctic CH₄ sources, in Table 1. Although we concentrate on the bottom-up budget in our discussion here, the top-down budget is not without issues. Top-down inverse modeling estimates for the Arctic are limited by relatively few atmospheric measurements in the Arctic [Bruhwiler *et al.*, 2014], tropospheric modeling capabilities [Houweling *et al.*, 1999], and uncertainty surrounding the hydroxyl radical, the primary atmospheric sink for CH₄ [Montzka *et al.*, 2011]. But top-down budgets are mass balanced by design, which is not the case for the bottom-up sums of independent studies.

2. Wetlands and Small Ponds

The past decade has seen increasing recognition of the importance of lakes and ponds as CH₄ sources in the Arctic and subarctic [Bastviken *et al.*, 2011; Walter *et al.*, 2006; Wik *et al.*, 2016b] and that those lakes and ponds' emissions are climate sensitive [Arp *et al.*, 2016; Tan and Zhuang, 2015; Thornton *et al.*, 2015]. Lakes and ponds are now seen as a distinct CH₄ source, apart from wetlands. The underlying methanogenic processes leading to CH₄ release from lakes and wetlands are similar, but not identical. For instance, the CH₄ flux from wetlands depends, in complex ways, on the water table [Brown *et al.*, 2014; Turetsky *et al.*, 2014], whereas more constant water in even small lakes eliminates this variable; further, lakes of various sizes can be active CH₄ producers throughout both their ice-free season and during winter [Walter *et al.*, 2008; Wik *et al.*, 2011], whereas wetlands may or may not be depending on seasonal weather. Many high-latitude lakes, for instance, have been found to release a substantial fraction of their annual CH₄ following ice-out in spring [Jammet *et al.*, 2015; Wik *et al.*, 2016b].

Table 1. Arctic CH₄ Budget; Bottom-Up Versus Top-Down^a

	Tgy ⁻¹	Study
<i>Bottom-Up Estimates</i>		
Lakes and ponds > 50°N	16.5 ± 9.2	Wik et al. [2016b]
Lakes and ponds > 60°N (bLake4Me model)	11.9	Tan and Zhuang [2015]
Rivers and streams > 54°N	0.3	Bastviken et al. [2011]
Rivers and streams > 54°N	7.5	Stanley et al. [2016]
Reservoirs > 54°N	1.2	Bastviken et al. [2011]
Arctic Ocean + Beaufort and Chukchi Seas (<82°N)	2	Kort et al. [2012]
ESAS	2.9	Thornton et al. [2016]
ESAS	17	Shakhova et al. [2014]
Wetlands > 60°N	23.2	Zhang et al. [2004]
Wetlands > 53.1°N (CarbonTracker prior model, based on Bergamaschi et al. [2005])	31	Bruhwiller et al. [2014]
Wetlands > 50°N (ORCHIDEE model)	31 ± 5	Bousquet et al. [2011]
Sources sum (minimum–maximum)	59.7 (36.9–89.4)	
<i>Top-Down Inverse Model Estimates</i>		
>60°N, all natural sources	23 ± 5	Bruhwiller et al. [2014] Saunois et al. [2016]
ESAS	0–4.5	Berchet et al. [2016]

^aRecent bottom-up estimates for various Arctic CH₄ source flux strengths are sorted into categories of lakes and ponds, rivers and streams, reservoirs, Arctic Ocean, ESAS, and wetlands. Estimates are based on extrapolations of measurements, except for the three process models noted. Note that the latitude bands differ, which partly account for the ultimate bottom-up uncertainty seen here. Arctic Ocean flux is from the reported 2 mg m⁻² d⁻¹ extrapolated over 10 × 10⁹ km² of seasonally ice-free Arctic Ocean regions for 100 ice-free days [Kort et al., 2012]. Rivers and streams high estimate is based on the Stanley et al. [2016] global fluvial flux database distributed into fluvial surface areas reported by Bastviken et al. [2011]. Sum uses averages of the all estimates per category. Minimum uses category low values and lower bound of the Wik et al. [2016b] lake estimates; maximum uses category high values and upper bounds of ORCHIDEE wetland model and the Wik et al. [2016b] lake estimates. Including subarctic and boreal wetlands from 45°N to 60°N would add 34 Tgy⁻¹ to the Zhang et al. [2004] wetland estimate.

Surprisingly, even uncertainties of total wetland area remain—not only in the Arctic [Bridgman et al., 2013]. Some of this uncertainty may arise from a conflation of wetlands with small ponds. The definition of wetlands may contribute to a double-counting problem. Widely used descriptions of wetlands (standing water ≤ 2–2.5 m deep) [Cowardin et al., 1979; Tiner et al., 2015] overlap with a vast array of high-latitude lakes and ponds, which tend to be small and can be less than 1 m deep—especially lakes and ponds in permafrost peatlands and thermokarst lakes [West and Plug, 2008]. Some wetland studies explicitly include shallow ponds [Melton et al., 2013]. Many shallow high-latitude lakes and ponds are persistent landscape features [Smith et al., 2007], unlike other seasonally inundated wetland areas (e.g., ecosystems adapted to flooded conditions) [Smith et al., 2007].

The error induced by double-counting of lake area emissions again as wetland area emissions is somewhat mollified by the fact that small lakes may have a greater annual emission potential than some small wetlands.

However, productive fens can closely resemble small, shallow ponds' CH₄ emission magnitude [Bubier et al., 1993; McEnroe et al., 2009; Pelletier et al., 2007]. It is precisely near the hazily defined wetland-or-lake boundary that the greatest CH₄ emissions per area are expected. The small water bodies which are most likely to be lumped into wetland areas cover an impressive expanse—lakes and ponds smaller than 0.1 km² cover in total about 1 × 10⁶ km² (equal to more than half the area of Alaska; Figure 1) [Verpoorter et al., 2014] —10% or more of the—uncertain—global wetland area [Melton et al., 2013].




Figure 1. Visualization of total area of small ponds which are most likely to be counted as wetlands in assessments. Verpoorter et al. [2014] estimate that such sized water bodies cover roughly 1 × 10⁶ km². In perspective this is more than half the area of the largest U.S. state, Alaska.




Figure 2. (a) Stordalen Mire, northern Sweden, 68°21'N, 19°02'E, at 2 m resolution. Worldview image (source: Digital Globe). Open lake waters are highlighted in blue (the large lake Torneträsk at the top of image is not highlighted). (b) Stordalen Mire at 50 m resolution, similar to resolution of lake database in [Verpoorter *et al.*, 2014], pixelated Figure 2a with lake pixel brightness tolerance of 10% from underlying open water pixel color. (c) Stordalen Mire at 350 m resolution, similar to resolution of lake database in [Lehner and Döll [2004], lake pixels calculated as in Figure 2b.

Models of regional and global CH₄ emissions utilizing lake databases with pixel sizes larger than extant lake sizes implicitly exclude these small water bodies. Due to previous limitations in remote sensing techniques, many wetland estimates have been based on lower resolution imagery [Lehner and Döll, 2004] and only exclude lakes < 0.1 km², such a lake is ~350 m in diameter. So far, only a very recent inventory, presented in the Global Water Bodies database (GLOWABO), accounts for smaller lakes, now as small as 0.002 km² [Verpoorter *et al.*, 2014]. Ironically, 0.002 km² is 40 times smaller than a common wetland/lake definition cutoff size, 0.08 km² [Tiner *et al.*, 2015], so the smallest lakes may, even if detected, sometimes be counted as wetlands. Figure 2 shows the effect of various pixel sizes on lake detection in a lake dense area in northern Sweden—a 49% loss of estimated water body area due to pixel size in this particular example; even GLOWABO misses many of the smallest ponds. Upscaling lake CH₄ emission estimates without including these small lakes not only excludes a large areal source but also excludes the strongest lake CH₄ source on a per unit area basis, as small water bodies are known to be hotspots of CH₄ production [Laurion *et al.*, 2010]. Conversely, a small wetland is not expected to be, per unit area, a larger CH₄ emitter than a large wetland. Additionally, it was recently shown that spatial and temporal sampling biases of CH₄ lake emissions are common and are likely to lead to underestimates of total lake emissions [Wik *et al.*, 2016a]. In the top-down total Arctic CH₄ budget, any such overlooked lake emissions due to sampling biases may have been counted as wetland emissions, but

increasingly accurate lake emission estimates suggest the need to include lake distributions, distinct from wetland distributions, as a specific input prior in top-down emission estimates. Specifically, in scaling bottom-up emission estimates, lake or wetland emissions are multiplied by lake or wetland area. If some amount of area is counted in both categories—from low-resolution maps or fuzzy definitions—double-counting of CH₄ emissions will exist in bottom-up estimates. Generally, models that attempt to avoid the double-counting problem are necessarily limited by the resolution of lake databases used for masking out lakes embedded in wetlands [Melton *et al.*, 2013].

Although wetlands are now reported separately from freshwater systems [Cole *et al.*, 2007] in global bottom-up and top-down accountings of CH₄ [Kirschke *et al.*, 2013], the separation currently appears to be more wishful than realistic. A recent bottom-up estimate that accounts for water bodies as small as 0.002 km² noted that lakes and ponds alone may account for as much as 70% of the inverse-modeling (top-down) $23 \pm 5 \text{ Tg yr}^{-1}$ predicted CH₄ emissions [Bruhwiler *et al.*, 2014] from all northern natural sources—sources which have been historically thought to be dominated by wetlands [Wik *et al.*, 2016b]. Wetlands are expected to dominate because nearly 50% of the world's wetlands lie between 50° and 70°N [Tiner *et al.*, 2015]. In our view, the reason for this discrepancy is a double-counting of the contribution of small lakes within wetlands to total CH₄ emissions—these small water bodies may be included in both lake and wetland bottom-up estimates. Similar double-counting problems also arise with streams and rivers embedded in wetland landscapes, as discussed below. Double-counting wetlands as lakes and vice-versa would always increase the bottom-up estimates of the natural sources in the CH₄ budget. In fact, the bottom-up estimate of natural sources is significantly higher than the top-down estimate [Kirschke *et al.*, 2013]. Improved resolution (smaller pixels) in remote sensing data of high-latitude landscape types [Drusch *et al.*, 2012; Jawak and Luis, 2013] may improve bottom-up estimates. Higher resolution will allow the discrimination of more small lakes embedded in wetlands, though regional calibration and ground truth studies will be required.

3. Sea Challenges

One of the surprises in CH₄ studies has been reports of substantial emissions to the atmosphere from shallow shelf seas. The ocean has long been regarded as only a small contributor to global CH₄ [Rhee *et al.*, 2009], but shallow seas are a different matter. Seafloor CH₄ sources—including bubble plumes—have a far better chance of reaching the atmosphere if the water is shallow, due to less time for the bubbles to dissolve, or for CH₄ to oxidize during upward advective transport [McGinnis *et al.*, 2006]. Small sea-air CH₄ fluxes were first suggested in the 1990s in the Beaufort Sea [Kvenvolden *et al.*, 1993], as well as moderate fluxes from the Beaufort and Chukchi Seas and Arctic Ocean [Kort *et al.*, 2012].

The largest shallow shelf sea in the world is the combined Laptev, East Siberian, and Chukchi Sea shelf (collectively, East Siberian Arctic Shelf, or ESAS), about $2.1 \times 10^6 \text{ km}^2$ with an average depth of 62 m [Jakobsson, 2002]. Understanding the scale of sea-air CH₄ fluxes from the ESAS has proven to be a challenge. Early reports of high fluxes from the Laptev and East Siberian Seas based on gas transfer models [Shakhova *et al.*, 2010] have been followed by even higher estimates based on counting bubbles from seabed gas plumes using sonar [Shakhova *et al.*, 2014]. CH₄ in shallow seas has been reported in other high-latitude locations [Kodovska *et al.*, 2016; Myhre *et al.*, 2016; Schmale *et al.*, 2010], though seawater CH₄ does not necessarily pre-empt substantial increases in atmospheric CH₄ [Myhre *et al.*, 2016]. There are a wide variety of potential sources of CH₄ in the waters of the ESAS. As the ESAS was above sea level at the last glaciation, it contains substantial subsea permafrost and organic material originally formed and frozen subaerially [Dmitrenko *et al.*, 2011]. Additionally, the Laptev and East Siberian Seas are strongly influenced by terrestrial organic carbon input from rivers, providing a modern source of C to the seas [Charkin *et al.*, 2011; Semiletov *et al.*, 2005]. Complicating matters further, the age of the carbon in the present day terrestrial organic matter source may be old or young, C released from thawing permafrost—or C in organic material produced in the annual cycle of plant growth. Coastal erosion of thawing permafrost shorelines provides yet another carbon input into the Arctic system [Lantuit *et al.*, 2013]. All of these marine and shore processes might contribute C to the Arctic CH₄ cycle.

To date, atmospheric observations cannot account for extremely high fluxes of CH₄ from the Laptev and East Siberian Seas (Table 1) [Berchet *et al.*, 2016], and recent reports have shown that not even all nearshore areas can be substantial sources [Overduin *et al.*, 2015] due to CH₄ oxidation in the sediment, and offshore seep sites can be limited in spatial extent [Thornton *et al.*, 2016]. Such heterogeneity is typical for all CH₄ sources, which complicates extrapolating regional total emissions from local measurements.

4. Isotopic Solutions

Lake, wetland, and sea emissions should be accounted for separately in CH₄ budgets, yet how can this be accomplished? Although variations in $\delta^{13}\text{C-CH}_4$ can help separate thermogenic and biogenic CH₄, and $\delta^{13}\text{C-CH}_4$ has been linked to methanogenic community changes along a thaw gradient in wetlands and




Figure 3. Ranges in site-specific mean $\delta^{13}\text{C-CH}_4$ reported from Arctic and subarctic lakes, wetlands, and shallow marine sediment from multiple data sources [Bouchard et al., 2015; Brosius et al., 2012; Coffin et al., 2013; Dove et al., 1999; Koch et al., 2009; Kuhlmann et al., 1998; McCalley et al., 2014; Nisbet, 2005; Overduin et al., 2015; Quay et al., 1988; Sapart et al., 2016; Sriskantharajah et al., 2012; Thompson et al., 2016; Walter et al., 2008; Wik, 2016]. Figure data are collected in the supporting information.

may thus shift global atmospheric $\delta^{13}\text{C-CH}_4$ in a warmer climate [McCalley et al., 2014], using this most common CH_4 isotope system for separating lakes, wetland, and shallow sea emissions is futile (Figure 3).

Recent work using a single isotope system suggests that the resumption of atmospheric CH_4 growth from 2007 onward is not due to new Arctic sources [Nisbet et al., 2016]. However, the stable isotopic signature of CH_4 long stored in subsea reservoirs, or produced from recently mobilized Pleistocene age C (from coastal erosion or terrestrial permafrost), may deceptively appear to match modern CH_4 biologically produced in wetlands if looking at only $\delta^{13}\text{C-CH}_4$ [Sapart et al., 2016].

Additional isotope systems, including $\Delta^{14}\text{C-CH}_4$ and $\delta\text{D-CH}_4$ may help in certain situations [Walter Anthony et al., 2012], but such measurements remain relatively uncommon in the literature, and the aforementioned remobilization of radiocarbon-depleted sources may inhibit the usefulness of $\Delta^{14}\text{C-CH}_4$ if old C is entering the modern carbon cycle. Regional assessments using multiple CH_4 isotopologues in the midlatitudes, however, have begun to appear [Townsend-Small et al., 2016]. Other techniques which are only now becoming available, such as monitoring exotic CH_4 isotopologues, e.g., $^{13}\text{CH}_3\text{D}$ and $^{12}\text{CH}_2\text{D}_2$ [Douglas et al., 2016], are promising for providing additional information, including the formation temperature of the CH_4 , and source and sink strengths [Whitehill et al., 2017], but presently are rarely applied due to methodological difficulty and cost. Ethane (C_2H_6) has been used to trace abiogenic CH_4 sources [Simpson et al., 2012] and has been occasionally used in characterizing high-latitude thermogenic CH_4 sources [Walter Anthony et al., 2012].

5. Rivers, Streams, and Reservoirs

Rivers are an additional CH_4 source in the Arctic [Kling et al., 1992; Striegl et al., 2012], and recognition that small rivers and streams may represent an additional uncounted CH_4 source has been growing in recent years [Crawford et al., 2014; McGinnis et al., 2016]. Five years ago, a data-limited synthesis estimated global CH_4 emissions from rivers at 1.5 Tg yr^{-1} [Bastviken et al., 2011]; a more recent synthesis reports 26.8 Tg yr^{-1} [Stanley et al., 2016]. The former report estimated fluvial CH_4 emissions from 54 to 66°N at 0.2 Tg yr^{-1} and

>66°N at 0.1 Tg yr⁻¹. The latter report included dozens of river and stream sites across the pan-Arctic, though a latitude-specific emission was not reported; we report in Table 1 an estimate based on fluvial area > 54°N and the new synthesis data from [Stanley *et al.*, 2016]. As with lakes, double-counting issues with resolution can occur; for example, care must be taken when counting rivers in wetlands regions. We are not aware of published isotopic studies of CH₄ emissions from streams and rivers at high latitudes; however, for reasons discussed above, we expect that one-dimensional isotopic studies will not yield meaningful discrimination of the relative importance of small rivers and streams to the CH₄ budget. Finally, reservoirs are sometimes collected separately from lakes, though their total influence at high latitudes is thought to be small [Bastviken *et al.*, 2011].

6. Conclusions

The picture or, rather, our appreciation of high-latitude CH₄ emissions from terrestrial aquatic sources seems to have grown much more complex. The Arctic CH₄ top-down inventory (23 ± 5 Tg yr⁻¹) [Bruhwiler *et al.*, 2014; Saunio *et al.*, 2016] and our rough bottom-up CH₄ inventory (59.7 (range 36.9–89.4) Tg yr⁻¹) do not match (Table 1). Although our inventory in Table 1 is hampered by variability in definitions of the pan-Arctic used in various studies, the mismatch remains. We do encourage future studies, as much as possible, to provide model and inventory outputs for a common set of latitude ranges (we suggest 50°N, >60°N, and above the Arctic Circle) to simplify intercomparisons.

Even 3 years ago, natural terrestrial CH₄ emission estimates were largely regarded as primarily wetlands [Kirschke *et al.*, 2013], with a global ratio of 217:40 for wetlands:inland waters, based on bottom-up estimates. A more recent synthesis changes that ratio to 185:122 [Saunio *et al.*, 2016], a dramatic change due to new knowledge of lake and pond, and river and streams emissions, but without concomitant changes of wetland total areas. In the last 3 years, estimates of global oceanic sources have actually been revised down, from 18 Tg yr⁻¹ to 14 Tg yr⁻¹, reflecting reduced uncertainty about the magnitude of shallow sea sources, though these sea sources certainly exist. High-latitude postglacial lakes that are permanent will likely increase CH₄ fluxes to the atmosphere in a warmer climate, independently of permafrost thaw [Thornton *et al.*, 2015]. Alongside that, large changes in the distribution and abundance of high-latitude thermokarst lakes are widely anticipated in the future [Andresen and Lougheed, 2015]. The challenges in interpreting the importance of CH₄ fluxes from any newly discovered or better-quantified sources were concisely stated in 2010 by Petrenko *et al.*: “A newly discovered CH₄ source is not necessarily a changing source, much less a source that is changing in response to Arctic warming.” [Petrenko *et al.*, 2010]. For top-down estimates, which provide a mass-balanced analysis of atmospheric observations, any newly discovered source may displace a portion of known sources in the budget—such as freshwater sources somewhat did recently for wetlands—or new sinks must be found.

Given the high prevalence of wetlands in boreal and Arctic zones, we vitally need ways to reliably discriminate CH₄ emissions from lakes, wetlands, reservoirs, streams and rivers, and shallow seas. δ¹³C-CH₄ studies alone are likely inadequate for this purpose, without a much greater understanding of what controls δ¹³C-CH₄ variability, beyond simply site type (Figure 3). There is evidence that studies of other CH₄ isotopologues may provide tools for such CH₄ source discrimination, but at this time, available data sets remain limited. Still, such tools are likely to benefit both top-down modeling and bottom-up in situ and model-based inventories of Arctic CH₄. It seems presumptuous to predict the effects of postulated future landscape changes (subaerial or subsea) on CH₄ fluxes to the atmosphere when we do not yet adequately separate wetlands from lakes and seas in the present and likely double-count sources in assessments. Improved landscape remote sensing at high latitudes (and ground truthing of that data in key regions), along with consistently applied definitions of wetlands that clearly distinguish wetland areas from lakes and ponds, are vital keys to reliable bottom-up CH₄ inventories at high latitudes.

Acknowledgments

Funding provided by Vetenskapsrådet, via the IZOMET-FS “Distinguishing Arctic CH₄ Sources to the Atmosphere Using Inverse Analysis of High Frequency CH₄, ¹³CH₄ and CH₃D Measurements” project grant to P.C. All data used in this study are either shown in the table or text, or are provided in the supplementary material.

References

- Andresen, C. G., and V. L. Lougheed (2015), Disappearing Arctic tundra ponds: Fine-scale analysis of surface hydrology in drained thaw lake basins over a 65 year period (1948–2013), *J. Geophys. Res. Biogeosci.*, 120, 466–479, doi:10.1002/2014JG002778.
- Arp, C. D., B. M. Jones, G. Grosse, A. C. Bondurant, V. E. Romanovsky, K. M. Hinkel, and A. D. Parsekian (2016), Threshold sensitivity of shallow Arctic lakes and sublake permafrost to changing winter climate, *Geophys. Res. Lett.*, 43, 6358–6365, doi:10.1002/2016gl068506.
- Bastviken, D., L. J. Tranvik, J. A. Downing, P. M. Crill, and A. Enrich-Prast (2011), Freshwater methane emissions offset the continental carbon sink, *Science*, 331(6013), 50–50, doi:10.1126/science.1196808.

- Berchet, A., et al. (2016), Atmospheric constraints on the methane emissions from the East Siberian Shelf, *Atmos. Chem. Phys.*, 16(6), 4147–4157, doi:10.5194/acp-16-4147-2016.
- Bergamaschi, P., M. Krol, F. Dentener, A. Vermeulen, F. Meinhardt, R. Graul, M. Ramonet, W. Peters, and E. J. Dlugokencky (2005), Inverse modelling of national and European CH₄ emissions using the atmospheric zoom model TM5, *Atmos. Chem. Phys.*, 5(9), 2431–2460, doi:10.5194/acp-5-2431-2005.
- Bouchard, F., I. Laurion, V. Preskienis, D. Fortier, X. Xu, and M. J. Whiticar (2015), Modern to millennium-old greenhouse gases emitted from ponds and lakes of the Eastern Canadian Arctic (Bylot Island, Nunavut), *Biogeosciences*, 12(23), 7279–7298, doi:10.5194/bg-12-7279-2015.
- Bousquet, P., et al. (2011), Source attribution of the changes in atmospheric methane for 2006–2008, *Atmos. Chem. Phys.*, 11(8), 3689–3700, doi:10.5194/acp-11-3689-2011.
- Bridgman, S. D., H. Cadillo-Quiroz, J. K. Keller, and Q. Zhuang (2013), Methane emissions from wetlands: biogeochemical, microbial, and modeling perspectives from local to global scales, *Glob. Change Biol.*, 19(5), 1325–1346, doi:10.1111/gcb.12131.
- Brosius, L. S., K. M. Walter Anthony, G. Grosse, J. P. Chanton, L. M. Farquharson, P. P. Overduin, and H. Meyer (2012), Using the deuterium isotope composition of permafrost meltwater to constrain thermokarst lake contributions to atmospheric CH₄ during the last deglaciation, *J. Geophys. Res.*, 117, G01022, doi:10.1029/2011JG001810.
- Brown, M. G., E. R. Humphreys, T. R. Moore, N. T. Roulet, and P. M. Lafleur (2014), Evidence for a nonmonotonic relationship between ecosystem-scale peatland methane emissions and water table depth, *J. Geophys. Res. Biogeosci.*, 119, 826–835, doi:10.1002/2013JG002576.
- Bruhwyler, L., E. Dlugokencky, K. Masarie, M. Ishizawa, A. Andrews, J. Miller, C. Sweeney, P. Tans, and D. Worthy (2014), CarbonTracker-CH₄: An assimilation system for estimating emissions of atmospheric methane, *Atmos. Chem. Phys.*, 14(16), 8269–8293, doi:10.5194/acp-14-8269-2014.
- Bubier, J. L., T. R. Moore, and N. T. Roulet (1993), Methane emissions from wetlands in the midboreal region of Northern Ontario, Canada, *Ecology*, 74(8), 2240–2254, doi:10.2307/1939577.
- Charkin, A. N., O. V. Dudarev, I. P. Semiletov, A. V. Kruhmalev, J. E. Vonk, L. Sánchez-García, E. Karlsson, and Ö. Gustafsson (2011), Seasonal and interannual variability of sedimentation and organic matter distribution in the Buor-Khaya Gulf: The primary recipient of input from Lena River and coastal erosion in the southeast Laptev Sea, *Biogeosciences*, 8(9), 2581–2594, doi:10.5194/bg-8-2581-2011.
- Coffin, R. B., J. P. Smith, R. E. Plummer, B. Yoza, R. K. Larsen, L. C. Millholland, and M. T. Montgomery (2013), Spatial variation in shallow sediment methane sources and cycling on the Alaskan Beaufort Sea Shelf/Slope, *Mar. Pet. Geol.*, 45, 186–197, doi:10.1016/j.marpetgeo.2013.05.002.
- Cole, J., et al. (2007), Plumbing the Global Carbon Cycle: Integrating Inland Waters into the Terrestrial Carbon Budget, *Ecosystems*, 10(1), 172–185, doi:10.1007/s10021-006-9013-8.
- Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe (1979), Classification of wetlands and deepwater habitats of the United States, *Report Rep. 79/31*, pp. 0–103, Washington, D. C.
- Crawford, J. T., E. H. Stanley, S. A. Spawn, J. C. Finlay, L. C. Loken, and R. G. Striegl (2014), Ebullitive methane emissions from oxygenated wetland streams, *Glob. Change Biol.*, 20(11), 3408–3422, doi:10.1111/gcb.12614.
- Dmitrenko, I. A., S. A. Kirillov, L. B. Tremblay, H. Kassens, O. A. Anisimov, S. A. Lavrov, S. O. Razumov, and M. N. Grigoriev (2011), Recent changes in shelf hydrography in the Siberian Arctic: Potential for subsea permafrost instability, *J. Geophys. Res.*, 116, C10027, doi:10.1029/2011JC007218.
- Douglas, P. M. J., et al. (2016), Diverse origins of Arctic and Subarctic methane point source emissions identified with multiply-substituted isotopologues, *Geochim. Cosmochim. Acta*, doi:10.1016/j.gca.2016.05.031.
- Dove, A., N. Roulet, P. Crill, J. Chanton, and R. Bourbonniere (1999), Methane dynamics of a northern boreal beaver pond, *Ecoscience*, 6(4), 577–586, doi:10.1080/11956860.1999.11682548.
- Drusch, M., et al. (2012), Sentinel-2: ESA's optical high-resolution mission for GMES operational services, *Remote Sens. Environ.*, 120, 25–36, doi:10.1016/j.rse.2011.11.026.
- Houweling, S., T. Kaminski, F. Dentener, J. Lelieveld, and M. Heimann (1999), Inverse modeling of methane sources and sinks using the adjoint of a global transport model, *J. Geophys. Res.*, 104(D21), 26,137–26,160, doi:10.1029/1999JD900428.
- Jakobsson, M. (2002), Hypsometry and volume of the Arctic Ocean and its constituent seas, *Geochem. Geophys. Geosyst.*, 3(5), 1–18, doi:10.1029/2001GC000302.
- Jammet, M., P. Crill, S. Dengel, and T. Friborg (2015), Large methane emissions from a subarctic lake during spring thaw: Mechanisms and landscape significance, *J. Geophys. Res. Biogeosci.*, 120, 2289–2305, doi:10.1002/2015JG003137.
- Jawak, S. D., and A. J. Luis (2013), A spectral index ratio-based Antarctic land-cover mapping using hyperspatial 8-band WorldView-2 imagery, *Polar Sci.*, 7(1), 18–38, doi:10.1016/j.polar.2012.12.002.
- Kirschke, S., et al. (2013), Three decades of global methane sources and sinks, *Nat. Geosci.*, 6(10), 813–823, doi:10.1038/ngeo1955.
- Kling, G., G. Kipphut, and M. Miller (1992), The flux of CO₂ and CH₄ from lakes and rivers in arctic Alaska, *Hydrobiologia*, 240(1–3), 23–36, doi:10.1007/bf00013449.
- Koch, K., C. Knoblauch, and D. Wagner (2009), Methanogenic community composition and anaerobic carbon turnover in submarine permafrost sediments of the Siberian Laptev Sea, *Environ. Microbiol.*, 11(3), 657–668, doi:10.1111/j.1462-2920.2008.01836.x.
- Kodovska, F. G.-T., K. J. Sparrow, S. A. Yvon-Lewis, A. Paytan, N. T. Dimova, A. Lecher, and J. D. Kessler (2016), Dissolved methane and carbon dioxide fluxes in Subarctic and Arctic regions: Assessing measurement techniques and spatial gradients, *Earth Planet. Sci. Lett.*, 436, 43–55, doi:10.1016/j.epsl.2015.12.002.
- Kort, E. A., et al. (2012), Atmospheric observations of Arctic Ocean methane emissions up to 82° north, *Nat. Geosci.*, 5(5), 318–321, doi:10.1038/ngeo1452.
- Kuhlmann, A. J., D. E. J. Worthy, N. B. A. Trivett, and I. Levin (1998), Methane emissions from a wetland region within the Hudson Bay Lowland: An atmospheric approach, *J. Geophys. Res.*, 103(D13), 16,009–16,016, doi:10.1029/98JD01024.
- Kvenvolden, K. A., M. D. Lilley, T. D. Lorenson, P. W. Barnes, and E. McLaughlin (1993), The Beaufort Sea continental shelf as a seasonal source of atmospheric methane, *Geophys. Res. Lett.*, 20(22), 2459–2462, doi:10.1029/93GL02727.
- Lantuit, H., P. P. Overduin, and S. Wetterich (2013), Recent progress regarding permafrost coasts, *Permafr. Periglac. Process.*, 24(2), 120–130, doi:10.1002/ppp.1777.
- Laurion, I., W. F. Vincent, S. MacIntyre, L. Retamal, C. Dupont, P. Francus, and R. Pienitz (2010), Variability in greenhouse gas emissions from permafrost thaw ponds, *Limnol. Oceanogr.*, 55(1), 115–133, doi:10.4319/lo.2010.55.1.0115.
- Lehner, B., and P. Döll (2004), Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296(1–4), 1–22, doi:10.1016/j.jhydrol.2004.03.028.
- McCalley, C. K., et al. (2014), Methane dynamics regulated by microbial community response to permafrost thaw, *Nature*, 514(7523), 478–481, doi:10.1038/nature13798.

- McEnroe, N. A., N. T. Roulet, T. R. Moore, and M. Garneau (2009), Do pool surface area and depth control CO₂ and CH₄ fluxes from an ombrotrophic raised bog, James Bay, Canada?, *J. Geophys. Res.*, *114*, G01001, doi:10.1029/2007JG000639.
- McGinnis, D. F., J. Greinert, Y. Artemov, S. E. Beaubien, and A. Wuest (2006), Fate of rising methane bubbles in stratified waters: How much methane reaches the atmosphere?, *J. Geophys. Res.*, *111*, C09007, doi:10.1029/2005JC003183.
- McGinnis, D. F., N. Bilsley, M. Schmidt, P. Fietzek, P. Bodmer, K. Premke, A. Lorke, and S. Flury (2016), Deconstructing methane emissions from a small Northern European River: Hydrodynamics and temperature as key drivers, *Environ. Sci. Technol.*, doi:10.1021/acs.est.6b03268.
- Melton, J. R., et al. (2013), Present state of global wetland extent and wetland methane modelling: Conclusions from a model inter-comparison project (WETCHIMP), *Biogeosciences*, *10*(2), 753–788, doi:10.5194/bg-10-753-2013.
- Montzka, S. A., M. Krol, E. Dlugokencky, B. Hall, P. Jöckel, and J. Lelieveld (2011), Small interannual variability of global atmospheric hydroxyl, *Science*, *331*(6013), 67–69, doi:10.1126/science.1197640.
- Myhre, C. L., et al. (2016), Extensive release of methane from Arctic seabed west of Svalbard during summer 2014 does not influence the atmosphere, *Geophys. Res. Lett.*, *43*, 4624–4631, doi:10.1002/2016GL068999.
- Nisbet, E. G. (2005), Meth-MonitEU: Methane monitoring in the European Union and Russia, Final Report, Section 6, European Commission, Brussels.
- Nisbet, E. G., et al. (2016), Rising atmospheric methane: 2007–2014 growth and isotopic shift, *Global Biogeochem. Cycles*, *30*, 1356–1370, doi:10.1002/2016GB005406.
- Overduin, P. P., S. Liebner, C. Knoblauch, F. Günther, S. Wetterich, L. Schirrmeister, H.-W. Hubberten, and M. N. Grigoriev (2015), Methane oxidation following submarine permafrost degradation: Measurements from a central Laptev Sea shelf borehole, *J. Geophys. Res. Biogeosci.*, *120*, 965–978, doi:10.1002/2014JG002862.
- Pelletier, L., T. R. Moore, N. T. Roulet, M. Garneau, and V. Beaulieu-Audy (2007), Methane fluxes from three peatlands in the La Grande Rivière watershed, James Bay lowland, Canada, *J. Geophys. Res.*, *112*, G01018, doi:10.1029/2006JG000216.
- Petrenko, V. V., D. M. Etheridge, R. F. Weiss, E. J. Brook, H. Schaefer, J. P. Severinghaus, A. M. Smith, D. Lowe, Q. Hua, and K. Riedel (2010), Methane from the East Siberian Arctic Shelf, *Science*, *329*(5996), 1146–1147, doi:10.1126/science.329.5996.1146-b.
- Prather, M. J., C. D. Holmes, and J. Hsu (2012), Reactive greenhouse gas scenarios: Systematic exploration of uncertainties and the role of atmospheric chemistry, *Geophys. Res. Lett.*, *39*, L09803, doi:10.1029/2012GL051440.
- Quay, P. D., S. L. King, J. M. Lansdown, and D. O. Wilbur (1988), Isotopic composition of methane released from wetlands: Implications for the increase in atmospheric methane, *Global Biogeochem. Cycles*, *2*(4), 385–397, doi:10.1029/GB002i004p00385.
- Rhee, T. S., A. J. Kettle, and M. O. Andreae (2009), Methane and nitrous oxide emissions from the ocean: A reassessment using basin-wide observations in the Atlantic, *J. Geophys. Res.*, *114*, D12304, doi:10.1029/2008JD011662.
- Sapart, C. J., et al. (2016), The origin of methane in the East Siberian Arctic Shelf unraveled with triple isotope analysis, *Biogeosci. Discuss.*, *2016*, 1–22, doi:10.5194/bg-2016-367.
- Saunio, M., et al. (2016), The global methane budget 2000–2012, *Earth Syst. Sci. Data*, *8*, 697–751, doi:10.5194/essd-8-697-2016.
- Schmale, O., J. Schneider von Deimling, W. Gülzow, G. Nausch, J. J. Waniek, and G. Rehder (2010), Distribution of methane in the water column of the Baltic Sea, *Geophys. Res. Lett.*, *37*, L12604, doi:10.1029/2010GL043115.
- Schuur, E. A. G., et al. (2015), Climate change and the permafrost carbon feedback, *Nature*, *520*(7546), 171–179, doi:10.1038/nature14338.
- Semiletov, I., O. Dudarev, V. Luchin, A. Charkin, K. H. Shin, and N. Tanaka (2005), The East Siberian sea as a transition zone between Pacific-derived waters and Arctic shelf waters, *Geophys. Res. Lett.*, *32*, L10614, doi:10.1029/2005GL022490.
- Shakhova, N., I. Semiletov, A. Salyuk, V. Yusupov, D. Kosmach, and O. Gustafsson (2010), Extensive methane venting to the atmosphere from sediments of the East Siberian Arctic Shelf, *Science*, *327*(5970), 1246–1250, doi:10.1126/science.1182221.
- Shakhova, N., et al. (2014), Ebullition and storm-induced methane release from the East Siberian Arctic Shelf, *Nat. Geosci.*, *7*(1), 64–70, doi:10.1038/ngeo2007.
- Simpson, I. J., M. P. Sulbaek Andersen, S. Meinardi, L. Bruhwiler, N. J. Blake, D. Helmig, F. S. Rowland, and D. R. Blake (2012), Long-term decline of global atmospheric ethane concentrations and implications for methane, *Nature*, *488*(7412), 490–494, doi:10.1038/nature11342.
- Smith, L. C., Y. Sheng, and G. M. MacDonald (2007), A first pan-Arctic assessment of the influence of glaciation, permafrost, topography and peatlands on northern hemisphere lake distribution, *Permafrost. Periglac. Process.*, *18*(2), 201–208, doi:10.1002/ppp.581.
- Sriskantharajah, S., R. E. Fisher, D. Lowry, T. Aalto, J. Hatakka, M. Aurela, T. Laurila, A. Lohila, E. Kuitunen, and E. G. Nisbet (2012), Stable carbon isotope signatures of methane from a Finnish subarctic wetland, *Tellus B*, *64*, 18818, doi:10.3402/tellusb.v64i0.18818.
- Stanley, E. H., N. J. Casson, S. T. Christel, J. T. Crawford, L. C. Loken, and S. K. Oliver (2016), The ecology of methane in streams and rivers: patterns, controls, and global significance, *Ecol. Monogr.*, *86*(2), 146–171, doi:10.1890/15-1027.
- Striegl, R. G., M. M. Dornblaser, C. P. McDonald, J. R. Rover, and E. G. Stets (2012), Carbon dioxide and methane emissions from the Yukon River system, *Global Biogeochem. Cycles*, *26*, GB0E05, doi:10.1029/2012GB004306.
- Tan, Z., and Q. Zhuang (2015), Arctic lakes are continuous methane sources to the atmosphere under warming conditions, *Environ. Res. Lett.*, *10*(5), 054016.
- Thompson, H. A., J. R. White, L. M. Pratt, and P. E. Sauer (2016), Spatial variation in flux, d13C and d2H of methane in a small Arctic lake with fringing wetland in western Greenland, *Biogeochemistry*, *131*(17), doi:10.1007/s10533-016-0261-1.
- Thornton, B. F., M. Wik, and P. M. Crill (2015), Climate-forced changes in available energy and methane bubbling from subarctic lakes, *Geophys. Res. Lett.*, *42*, 1936–1942, doi:10.1002/2015GL063189.
- Thornton, B. F., M. C. Geibel, P. M. Crill, C. Humborg, and C.-M. Mörh (2016), Methane fluxes from the sea to the atmosphere across the Siberian shelf seas, *Geophys. Res. Lett.*, *43*, 5869–5877, doi:10.1002/2016GL068977.
- Tiner, R. W., M. W. Lang, and V. V. Klemas (2015), *Remote Sensing of Wetlands: Applications and Advances*, 574 pp., CRC Press, Boca Raton, Fla.
- Townsend-Small, A., et al. (2016), Using stable isotopes of hydrogen to quantify biogenic and thermogenic atmospheric methane sources: A case study from the Colorado Front Range, *Geophys. Res. Lett.*, *43*, 11,462–11,471, doi:10.1002/2016GL071438.
- Turetsky, M. R., et al. (2014), A synthesis of methane emissions from 71 northern, temperate, and subtropical wetlands, *Glob. Change Biol.*, *20*(7), 2183–2197, doi:10.1111/gcb.12580.
- Walter Anthony, K. M., P. Anthony, G. Grosse, and J. Chanton (2012), Geologic methane seeps along boundaries of Arctic permafrost thaw and melting glaciers, *Nat. Geosci.*, *5*(6), 419–426, doi:10.1038/ngeo1480.
- Walter, K. M., S. A. Zimov, J. P. Chanton, D. Verbyla, and F. S. Chapin (2006), Methane bubbling from Siberian thaw lakes as a positive feedback to climate warming, *Nature*, *443*(7107), 71–75, doi:10.1038/nature05040.
- Walter, K. M., J. P. Chanton, F. S. Chapin, E. A. G. Schuur, and S. A. Zimov (2008), Methane production and bubble emissions from arctic lakes: Isotopic implications for source pathways and ages, *J. Geophys. Res.*, *113*, G00A8, doi:10.1029/2007JG000569.
- Verpoorter, C., T. Kutser, D. A. Seekell, and L. J. Tranvik (2014), A global inventory of lakes based on high-resolution satellite imagery, *Geophys. Res. Lett.*, *41*, 6396–6402, doi:10.1002/2014GL060641.

- West, J. J., and L. J. Plug (2008), Time-dependent morphology of thaw lakes and taliks in deep and shallow ground ice, *J. Geophys. Res.*, *113*, F01009, doi:10.1029/2006JF000696.
- Whitehill, A. R., L. M. T. Joelsson, J. A. Schmidt, D. T. Wang, M. S. Johnson, and S. Ono (2017), Clumped isotope effects during OH and Cl oxidation of methane, *Geochim. Cosmochim. Acta*, *196*, 307–325, doi:10.1016/j.gca.2016.09.012.
- Wik, M. (2016), Emission of methane from northern lakes and ponds, PhD Thesis, Stockholm Univ., Stockholm.
- Wik, M., P. M. Crill, D. Bastviken, A. Danielsson, and E. Norback (2011), Bubbles trapped in arctic lake ice: Potential implications for methane emissions, *J. Geophys. Res.*, *116*, G03044, doi:10.1029/2011JG001761.
- Wik, M., B. F. Thornton, D. Bastviken, J. Uhlbäck, and P. M. Crill (2016a), Biased sampling of methane release from northern lakes: A problem for extrapolation, *Geophys. Res. Lett.*, *43*, 1256–1262, doi:10.1002/2015GL066501.
- Wik, M., R. K. Varner, K. W. Anthony, S. MacIntyre, and D. Bastviken (2016b), Climate-sensitive northern lakes and ponds are critical components of methane release, *Nat. Geosci.*, *9*(2), 99–105, doi:10.1038/ngeo2578.
- Vonk, J. E., et al. (2013), High biolability of ancient permafrost carbon upon thaw, *Geophys. Res. Lett.*, *40*, 2689–2693, doi:10.1002/grl.50348.
- Zhang, G. L., J. Zhang, Y. B. Kang, and S. M. Liu (2004), Distributions and fluxes of methane in the East China Sea and the Yellow Sea in spring, *J. Geophys. Res.*, *109*, C07011, doi:10.1029/2004JC002268.